



Lawns

Planting: Rake into the soil 2 level tablespoons per square metre prior to sowing seed.

Topdressing: Apply Multigro at 1½ level tablespoons per square metre early in August and a further ¾ level tablespoon per square metre in February.

In addition, apply 1¼ level tablespoons per square metre of Incitec Pivot GRAN-AM (Sulfate of Ammonia) late in September.

Turf: When establishing lawn using sod or turf, apply 1½ level tablespoons per square metre before laying turf. Water or work into the soil. When the turf is laid and after the first watering, give an initial topdressing at ¾ level tablespoon per square metre. Then move onto the full topdressing programme.

Measurements

1 level tablespoon of Multigro weighs approximately 30 grams. A 2 litre icecream container holds about 1.75 - 2 kilograms. 1 cup (250mL) will hold approximately 250 grams of Multigro.

A 25 or 40kg bag of Multigro is small enough to move around and stores conveniently out of the way, yet large enough to give you the advantage of a quantity buy.

NOTE: Adequate watering, disease and insect control are important in getting best results.



For more information on Multigro, please contact your local Incitec Pivot Distributor. Freecall 1800 009 832

www.incitecpivot.com.au

Incitec Pivot Limited ABN 42 004 080 264 Multigro and GRAN-AM are registered trademarks of Incitec Pivot Limited.



The multi-purpose garden fertiliser.



Multigro® – the pick of the bunch.

Feed them up on Multigro, the multipurpose plant growth booster.

Each big economical pack contains the major plant food nutrients in carefully balanced proportions, so your garden gets exactly what it needs.

Surround yourself with lush lawns, bountiful fruit trees, and healthy vegetables, flowers and shrubs. For rates and application refer to the recommendations within this brochure.

Multigro, the complete fertiliser for the complete garden.

Multigro Analysis:	N	P	K	S	Ca
	13.1%	4.5%	7.2%	15.4%	2.4%

Flowers and Ornamentals

Large Trees & Shrubs

At Planting: 3 level tablespoons worked into the soil to a depth of 20cm over a circular area 60cm across.

Side-Dressing: Side-dress in August and again in February with $\frac{3}{4}$ full cup per 5cm of trunk diameter measured 30cm above the ground. Spread under the canopy but keep away from the trunk.

Hibiscus

The same rates as for large trees and shrubs may be used. But hibiscus growers may find it more desirable to fertilise each month from August to May inclusive with 2 tablespoons per application.

Annuals

In Rows: $1\frac{1}{2}$ level tablespoons per metre of row worked into the soil before planting.

In Beds: 3 level tablespoons per square metre worked into the soil before planting.

Roses

At Planting: 3 level tablespoons per square metre of bed, worked well into the soil to a depth of 10cm.

Side-Dressing: $1\frac{1}{2}$ level tablespoons spread around the bush in August followed by a further $\frac{3}{4}$ level tablespoon when buds are forming.

Camellias, Azaleas etc

Multigro® is suitable for camellias and azaleas. Spread under and slightly beyond the canopy using $1\frac{1}{2}$ level tablespoons per square metre. Apply in late August and again in February.

Bulbs and Corn

3 level tablespoons per square metre of bed worked into the soil three weeks before planting. In perennial beds, work in a similar quantity when the soil is worked in the winter.

Vegetables

Root Crops

Potatoes: 3 level tablespoons per metre of row worked into the soil in the bottom of the row.

Carrots, Parsnip, Beetroot, Turnip etc: $1\frac{1}{2}$ level tablespoons per metre of row worked into the soil beside and below the seed.

Leafy Crops

Cabbage, Cauliflower, Brussel Sprouts, Broccoli, Silver Beet, Spinach, Lettuce, Rhubarb: 3 level tablespoons per square metre worked into the soil before planting. Then side-dress with 2 level tablespoons of Incitec Pivot GRAN-AM® (Sulfate of Ammonia) per metre of row half way through the growing period.

Sweet Corn

$1\frac{1}{2}$ level tablespoons per metre of row worked into the soil beside and below the seed. When tassels appear on top of plants, side-dress with 1 level tablespoon per metre of Incitec Pivot GRAN-AM (Sulfate of Ammonia).

Beans and Peas

$1\frac{1}{2}$ level tablespoons per metre of row worked into the soil beside and below the seed. After 3 weeks side-dress at half this rate.

Tomatoes

$1\frac{1}{2}$ level tablespoons per plant worked into the soil around the plant. After 5 weeks repeat the application as a side-dressing.

Melons, Pumpkins, Squash etc.

3 level tablespoons per hill worked into the soil before planting.

Fruit

General

Treat in very early spring (preferably August). Spread fertiliser evenly over the drip area, (the area directly under the canopy), but keep 30cm away from the trunk.

Apples and Pears

Younger Trees: $\frac{1}{2}$ cupful for each year of age up to 3 full cups.

Mature Trees: 3 to 5 full cups depending on size.

Peaches, Apricots, Plums, Nectarines, Cherries

Younger Trees: $\frac{3}{4}$ cupful for each year of age up to $4\frac{1}{2}$ full cups.

Mature Trees: $4\frac{1}{2}$ to 7 full cups depending on size.

Citrus

Younger Trees: $\frac{1}{2}$ cupful per year of age up to 4 full cups.

Mature Trees: 4 to 7 full cups depending on size. Apply in August and repeat the application in February at half rates.

Grape Vines

$\frac{3}{4}$ full cup per vine in August.

Bananas

Apply $\frac{1}{2}$ cupful per stool every ten weeks, (five times per year).

Strawberries

3 level tablespoons per metre of row, placed in two bands 10cm apart and 10cm deep. Plant between the bands. Side-dress with similar quantity at flowering stage.

