

Boost spring feed with strategic nitrogen

Clint Sheather
IPF Technical Agronomist

This winter has been wet and cool in many areas reducing winter feed growth. So, livestock producers may consider applying nitrogen to increase pasture growth to avoid a feed gap or boost silage or hay yields.

In southern Australia, pasture growth is often negligible in winter and limited in early spring. For example, perennial ryegrass is most productive at ambient temperatures of 18-25°C while soil temperatures below 10°C also minimise growth. So, spring can be a period where the demand for feed exceeds pasture growth.

Strategic applications of nitrogen at key times to manage feed deficits are more efficient than a fixed rate of nitrogen. Before applying nitrogen there are some things to consider:

- Is the pasture actively growing?
- Can excess growth be conserved as hay or silage?
- Are there other nutrients or soil constraints that will limit pasture growth?
- What is the return from the additional feed grown?

ECONOMICS OF N APPLICATION

The IPF [Feed Cost Calculator](#) can help determine the return on nitrogen investment – which will be influenced by moisture, pasture composition and density, soil fertility, grazing management, application rate and fertiliser product.

Trial work at Howlong in 2020, showed applications of both 30 and 60 kg/ha nitrogen produced more dry matter than the control (Figure 1).




Figure 1: Nitrogen response (kg DM per kg N) of Horizon phalaris in a dryland trial at Howlong. Source: Barenburg Seeds & Incitec Pivot Fertilisers 2020.



But an early spring application of Green Urea NV®, at both 30 and 60 kg/ha of nitrogen, produced significantly more dry matter than urea.

When the cost per tonne of dry matter was calculated (Table 1), even at \$1000+ per tonne, urea and Green Urea NV are still an economic way to produce additional feed.

In early spring, when the potential for nitrogen loss is greater, the additional investment in Green Urea NV of \$40/tonne produced more feed at a lower cost. Green Urea NV also provides additional logistical flexibility.

Table 1: Cost per tonne of dry matter based on a urea cost of \$1,200/t and Green Urea NV cost of \$1,240/t.

Timing	Winter	Early spring	Late spring
30 kg/ha of N as Urea	\$428	\$203	\$260
30 kg/ha of N as Green Urea NV	\$395	\$164	\$262
60 kg/ha of N as Urea	\$498	\$203	\$349
60 kg/ha of N as Green Urea NV	\$419	\$169	\$354

Boost spring feed with strategic nitrogen applications

RIGHT PRODUCT

Urea 46% N w/w

Urea is the most widely used high analysis granular nitrogen fertiliser. However, spring applications are prone to ammonia loss through volatilisation. Losses of up to 30% have been recorded where:

- limited moisture has dissolved but not moved the urea into the soil
- the soil temperature is over 15°C
- daytime temperatures are warm to hot
- the soil surface is drying out
- moderate to strong wind conditions are likely at the soil surface

Green Urea NV® 46% N w/w

Green Urea NV is granular urea coated with the urease inhibitor NBPT (N-(n-Butyl)-thiophosphoric triamide) to protect against volatilisation losses.

In spring, as temperatures increase, 20 to 30% of N applied to pastures can be lost through volatilisation. Green Urea NV can reduce potential volatilisation loss by 70%. Green Urea NV inhibits the activity of the urease enzyme, delaying the hydrolysis process and extending the time for incorporating before rainfall for a period of up to 14 days. This allows more time for the fertiliser to be safely spread or incorporated into the soil by rainfall or irrigation, where it can be used by pastures without delaying dry matter production. This also means a larger area can be treated ahead of each rain event. The wider application window reduces the risk that an application will be missed.

EASY N® 42.5% w/w

Easy N® is a high analysis liquid fertiliser present as urea ammonium nitrate (urea 50%, 25% ammonium, and 25% nitrate-nitrogen).

Easy N offers several advantages compared with urea. The presence of nitrate-nitrogen means a faster response on cold soils. It can be applied by fertigation or surface application through streaming and flat fan nozzles.

Application of Easy N to pastures should occur immediately after grazing to maximise dry matter production and avoid leaf burn to new growth. Easy N is still subject to volatilisation losses so a rain event of 10 mm or more is required to minimise nitrogen losses.

A minimum of 50 L/ha should be applied to actively growing pasture using flat fan nozzles while streaming nozzles should be considered when applying >50 L/ha of Easy N. Table 2 shows the rate comparison between urea / Green Urea NV and Easy N.

Table 2: Comparison of the rate of Urea and Easy N required to achieve different N rates.

Nitrogen rate kg/ha	Urea / Green Urea rate (46% N w/w)	EASY N rate (42.5% N w/w)
25	54	60
50	109	119
75	163	179
100	217	238
125	272	298
150	326	357
175	380	417
200	435	476

RIGHT TIMING

Pastures are most nitrogen deficient from late autumn to early spring when less nitrogen is being mineralised by the soil. Strategic applications in late autumn, before soil temperatures fall can provide additional winter feed. Similarly, late winter to early spring applications can produce additional feed for spring if supported by the right conditions.

Nitrogen applied soon after grazing is more efficient than an application onto a regrown pasture. Nitrogen responses decrease by 1% for every day the application is delayed, i.e., delaying nitrogen application for 10 days after the sheep or cattle come out of the paddock will reduce the potential nitrogen response by 10%.

RIGHT RATE

The best dry matter response is generally to applications of between 25 and 50 kg N/ha. In this range, dry matter will accumulate as nitrogen rate increases. However, beyond 50 kg N/ha, there is a point at which a positive economic outcome is unlikely. Each individual needs to consider their situation, input costs, response rate and conversion rate, to find that point.

Boost spring feed with strategic nitrogen applications

The response curve (Figure 2) shows that low rates of nitrogen are often not sufficient to provide a response while high rates are not used as efficiently and don't always produce more dry matter. Excessive rates can lead to losses through leaching, denitrification and volatilisation.




Figure 2: The pasture response to N fertiliser, both as the percentage of the maximum growth rate that can be achieved, and as the daily growth rate change for a single grazing rotation for a well fertilised perennial ryegrass pasture in late winter. Source: Eckard, 2010.

RIGHT AGRONOMY

Pasture condition

Pasture condition is key to the success of nitrogen applications. The pasture must be actively growing for nitrogen to produce additional dry matter. If the pasture is moisture stressed, the response to nitrogen will be significantly reduced. As a rule, a minimum of 50 mm of plant-available water is needed to support nitrogen responses.

Make sure there aren't other factors that will limit growth, such as phosphorus (Olsen P <15 mg/kg Colwell P <30 mg/kg 0-10 cm), sulphur (KCl40 <8 mg/kg 0-10 cm), or acidity.

Animal health

Nitrate poisoning in livestock can occur after applying high rates of nitrogen fertilisers – particularly combined with low temperatures and periods of moisture stress. Nitrate poisoning can also be related to pasture composition. It is recommended that grazing should not occur for 21 days after nitrogen applications.

REFERENCES

Eckard R (2010). Nitrogen-growth promotant for pastures.

Eckard RJ, & Franks DR (1998). Strategic nitrogen fertiliser use on perennial ryegrass and white clover pasture in north-western Tasmania. *Australian Journal of Experimental Agriculture*, 38(2), 155-160.

Smith AP, Christie KM, Rawnsley RP, & Eckard RJ (2018). Fertiliser strategies for improving nitrogen use efficiency in grazed dairy pastures. *Agricultural Systems*, 165, 274-282.

Suter H, Sultana H, Turner D, Davies R, Walker C, & Chen D (2013). Influence of urea fertiliser formulation, urease inhibitor and season on ammonia loss from ryegrass. *Nutrient cycling in agroecosystems*, 95(2), 175-185.

FURTHER INFORMATION

For more information or advice about staying on boosting spring pasture growth, feel free to contact me on 0475 439 316 or clint.sheather@incitecpivot.com.au.

You can also contact:

Lee Menhenett lee.menhenett@incitecpivot.com.au

Fiona McDonald fiona.mcdonald@incitecpivot.com.au

Jim Laycock jim.laycock@incitecpivot.com.au

ACKNOWLEDGMENTS

Incitec Pivot Fertilisers would like to acknowledge Barenburg Seeds for their contribution during the Howling nitrogen response trial in 2020.

DISCLAIMER

This is a guide only, which we hope you find useful as a general tool. While Incitec Pivot Fertilisers has taken all reasonable care in the preparation of this guide, it should not be relied on as a substitute for tailored professional advice and Incitec Pivot Fertilisers accepts no liability in connection with this guide.

Incitec Pivot Fertilisers manufactures and sources fertilisers from other suppliers. The fertiliser supply chain extends beyond the company's direct control, both overseas and within Australia. Incitec Pivot Fertilisers hereby expressly disclaims liability to any person, property or thing in respect of any of the consequences of anything done or omitted to be done by any person in reliance, whether wholly or in part, upon the whole or any part of the contents of this article.